

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/151479>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

© 2021 Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>.



Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

SARS-CoV-2 incidence and vaccine escape

AUTHORS

R.N. Thompson^{1,2,3,*}, E.M. Hill^{1,2,3,4}, J.R. Gog^{3,5}

*Correspondence to: robin.n.thompson@warwick.ac.uk

AFFILIATIONS

¹Mathematics Institute, University of Warwick, Coventry, UK

²Zeeman Institute for Systems Biology and Infectious Disease Epidemiology Research,
University of Warwick, Coventry, UK

³JUNIPER – Joint Universities Pandemic and Epidemiological Research,
<https://maths.org/juniper/>

⁴School of Life Sciences, University of Warwick, Coventry, UK

⁵Department of Applied Mathematics and Theoretical Physics, University of Cambridge,
Cambridge, UK

A recent Editorial¹ described the potential for the evolution of SARS-CoV-2 variants that render vaccines less effective (vaccine escape), assisted by waning immunity following vaccination. This raises a crucial question: how can COVID-19 exit strategies be planned with only a low vaccine escape risk?

A key component of any plausible strategy towards the permanent removal of non-pharmaceutical interventions (NPIs) involves ensuring low case numbers in the short-to-mid-term using NPIs and vaccination. Assuming a fixed vaccine escape mutation probability per infection (p), the risk of a vaccine escape variant arising in a specified time period is

$$\text{Prob}(\text{vaccine escape}) = 1 - (1 - p)^N,$$

where N represents the number of cases in that period.

Crucially, this indicates that the vaccine escape risk is sensitive to background incidence: the risk of an escape variant appearing within a fixed time is an increasing function of incidence (Fig 1). Reducing cases is not only beneficial for decreasing the pressure on healthcare systems, but also for lowering the vaccine escape risk.

Of course, there are fundamental differences between using NPIs and vaccines to lower incidence. When considering vaccines that do not prevent transmission entirely, there is an interplay between reduced cases at the population-level and the potential for selection for vaccine escape variants in infected vaccinated hosts²⁻⁴. A related question is whether it is most beneficial to vaccinate many individuals using single vaccine doses or fewer individuals twice. “Dose-sparing” strategies could in theory lead to selection for vaccine escape variants⁵. However, current evidence suggests tentatively that the net vaccine escape risk is lower when more hosts are vaccinated with single doses, due to reduced cases².




Figure 1. The risk that at least one vaccine escape variant arises in a time period of length t , for different daily numbers of cases. In this figure, the per infection probability of vaccine escape is $p = 2 \times 10^{-7}$ (for details, see Supplementary Material).

60

61 Despite its simplicity, our quantitative illustration demonstrates that strategies for
62 mitigating the vaccine escape risk should be explored. Reducing case numbers locally
63 should only be one component of these strategies. Travel restrictions to reduce the risk
64 of importing novel variants should be considered. We recognise that assessing the
65 escape variant emergence risk not only requires the variant to arise via mutation as
66 considered here, but also to grow to appreciable frequencies. This is a stochastic process,
67 depending on the availability of hosts to infect and the escape variant's fitness. A
68 reduction in cases leads to both a reduction in the risk of escape variants appearing and
69 a reduction in their subsequent establishment through transmission around the
70 population. Acquisition of additional mutations that are beneficial for the virus is also more
71 likely to be suppressed if incidence rates are reduced.

72

73 In summary, high SARS-CoV-2 incidence rates act to increase the vaccine escape risk.
74 Maintaining low case numbers using NPIs and vaccines is critical at the current time.

75

76

REFERENCES

77

78 1. The Lancet Infectious Diseases. An exceptional vaccination policy in exceptional
79 circumstances. *Lancet Infect. Dis.* **21**, 149 (2020).

80 2. Cobey, S., Larremore, D. B., Grad, Y. H. & Lipsitch, M. *Concerns about SARS-*
81 *CoV-2 evolution should not hold back efforts to expand vaccination.*
82 <https://nrs.harvard.edu/URN-3:HUL.INSTREPOS:37366988> (2021).

83 3. Saad-Roy, C. M., Morris, S. E., Metcalf, C. J. E., Mina, M. J., Baker, R. E., Farrar,
84 J., Holmes, E. C., Pybus, O. G., Graham, A. L., Levin, S. A., Grenfell, B. T. &
85 Wagner, C. E. Epidemiological and evolutionary considerations of SARS-CoV-2
86 vaccine dosing regimes. *Science*. eabg8663 (2021).

87 4. Gog, J. R., Hill, E. M., Danon, L. & Thompson, R. N. Vaccine escape in
88 heterogeneous populations: insights for SARS-CoV-2 from a simple model.
89 *medRxiv* (2021).

90 5. Bieniasz, P. D. The case against delaying Severe Acute Respiratory Syndrome
91 Coronavirus 2 (SARS-CoV-2) mRNA vaccine boosting doses. *Clin. Infect. Dis.*
92 ciab070 (2021).

93